

A Study on the Association
between Poverty,
Demographics, Family
Support, and Cancer Care
in Jharkhand, India

**Imteyaz Ahmad, Padmaja Gadiraju,
Punam Singh, Ranjit Kumar
Dehury, C Vanlalhruaii,
Parthsarathi Dehury**

Outline

Introduction

Literature Review

Objectives

Methodology

Results

Discussion & Conclusion

Introduction

- Cancer is a leading cause of morbidity and mortality worldwide. The burden of incidence, morbidity, and mortality disproportionately affects the developing world.
- The world is hampered by cancers attributable to infectious diseases and risks associated with diet, tobacco, alcohol, lack of exercise, and industrial exposures(Kanavos, 2006).
- India, marked by its diversity and complexity, is confronted with a pressing public health issue involving poverty and cancer care.
- Jharkhand, a state in eastern India, struggles with a multifaceted challenge concerning cancer care, where poverty plays a pivotal role in determining access to quality healthcare.

Literature Review

- A higher incidence of several cancers and lower survival rates are common among the poor. Poverty contributes to an increase in cancer incidence and mortality. Several factors are responsible for the increased mortality and morbidity from cancer among low-income people (Wilkes et al., 1994; Ratcliff et al., 2017; Callahan & Brintzenhofeszoc, 2015).
- A study found the effect of poverty on increasing the death rate of cancer disease. Low-income countries have a higher rate of cancer mortality than high-income countries. The GDP income significantly impacts individuals' health, quality of life, and cancer mortality rates (Alzu'Bi., et al., 2022).
- Many cancer cases in India are associated with tobacco use, infections, and other avoidable causes. The number of new cases of cancer diagnosed every year and the number of deaths is increasing in India (Mallath et al., 2014; Dhillonet al., 2018).
- A study found that lack of social support was associated with a higher cancer incidence and mortality risk. The study also identified that social support affects cancer onset and prognosis via a range of factors, including healthier lifestyles and adherence to therapeutic regimens (Ikeda et al., 2013)

Literature Review

- Indian oncology clinicians identified three key challenges in cancer care: practical constraints, cultural values, and structural conditions. The practical constraints include access and treatment. Cultural values include communication, stigma, and the clinic. Structural conditions include inequalities related to place, gender, and class (Broom & Doron, 2012).
- A study shows the age-standardized prevalence of cancer in India is estimated to be 97 per 100,000 persons, with a greater majority in urban areas. The average out-of-pocket spending on inpatient care in private facilities is three times greater than in public facilities (Rajpal et al., 2018).
- Public expenditure on cancer in India is low compared to high-income countries. Out-of-pocket payments account for most of the cancer expenditures, leading to catastrophic expenditures for patients and their families (Pramesh et al., 2014).
- A study found over 8,00,000 new cases and 5,50,000 deaths occur annually due to cancer in India, with the rural and underprivileged population representing the majority of patients (Das & Patro, 2010).

Objective

S

The complex relationship between cancer care and poverty was examined in this paper. The study identified the association of various demographic factors with cancer care. The study also analyses the family's role and the support network in cancer care.

Methodology

Mixed-method approach

Cross-sectional exploratory design

Structured interview schedule

Case study

Methodology

- The study is provided with ethical approval from the Institutional Review Board (IRB) at the University of Hyderabad.
- The interview schedule comprised sections on demography, access to cancer care facilities and socioeconomic conditions of cancer patients.
- The survey method allowed for the collection of quantitative data to analyse various aspects of cancer care. Case study methods explored the impact of social determinants, poverty and other factors.
- The target population for the survey includes cancer patients who are currently receiving treatment or have received treatment in Jharkhand in the last year.
- Ranchi Cancer Hospital and Research Centre was approached for quantitative data collection. A purposive sampling technique was used to collect data.

Methodology

- Cancer cases for in-depth-interviews were recruited from Ranchi Cancer Hospital and Research Centre and Rajendra Institute of Medical Sciences, Ranchi.
- The sample size was determined using the prevalence of cancer in Jharkhand. The sample calculated was 204 respondents.
- Three eligible adult participants were found in the oncology centres at different hospitals in Jharkhand; among them, one participant has undergone cancer treatment while two others were currently undergoing treatment.
- Descriptive statistics are generated as a part of the study findings using SPSS version 25.
- Cancer patients' lived experiences are represented in the case studies. The three cases highlight three major issues pertaining to the cancer impact; the first is poverty, the second is fear and family support, and the third is about social stigma.

Results

Socio-demographic
characteristics

Access to cancer care

Poverty and cancer

Fear and family support

Stigma and social
support

Socio-demographic characteristics

Respondents' characteristics	Category	Frequency	Percentage	Total (N)
Age	20 to 39 years	56	27.5	204
	40-59 years	123	60.3	
	60 years and above	25	12.3	
	Mean age	45.27		
Gender	Male	52	25.5	204
	Female	152	74.5	
Religion	Hindu	142	69.6	204
	Muslim	49	24.0	
	Christian	13	6.4	
Education	Not attended school	88	43.1	204
	Up to 10 th standard	78	38.2	
	10 th and above	38	18.6	
Type of Residence area	Urban	84	41.2	204
	Rural	87	42.6	
	Semi-Urban	33	16.2	
Infrastructure	Kutchha	30	14.7	204
	Semi-Pucca	94	46.1	
	Pucca	80	39.2	
Total Annual income of the family	Up to 100,000 INR	68	33.3	204
	100001-200000 INR	72	35.3	
	200001 INR and above	64	31.4	

Access to cancer care

	Respondents' characteristic	Treatment	Vaccinated (HPV)	Physiotherapy	Counselling services	Total
Age	20 to 39 years	46.4	1.8**	35.7*	66.1***	56
	40-59 years	48.8	17.9	51.2	35.0	123
	60 years and above	40.0	12.7	32.0	44.0	25
Gender	Male	42.3	17.3	38.5	61.5***	52
	Female	48.7	11.2	46.7	38.8	152
Religion	Hindu	45.8**	12.0	45.1***	40.8**	142
	Muslim	40.8	14.3	55.1	59.2	49
	Christian	84.6	15.4	0.00	30.8	13
Education	Not attended school	28.4***	10.2	44.3	42.0	88
	Up to 10 th standard	60.3	12.8	42.3	44.9	78
	11 th and above	63.2	18.4	50.0	50.0	38
Household income	Up to 100,000 INR	44.1	14.7	35.3**	47.1	68
	100001-200000 INR	45.8	11.1	56.9	41.7	72
	200001 INR and above	51.6	12.5	40.6	45.3	64
	Total	47.1	12.7	44.6	44.6	204
Significance: ***p<0.01, **p<0.05, *p<0.1						

Poverty and Cancer

- Mohammad Hushen (Name changed), a 38-year-old daily wage labourer, was battling with tongue and skin cancer. Hushen was not very educated. In his family, five members were there in aggregate (a couple and three children). Children were below seven years of age. Hushen was the sole earning member of the family. Four months ago, he was diagnosed with cancer. Despite being aware of the disease, he continued working as a daily wage labourer. He was more concerned about family expenses, including his treatment costs. His pain aggravated, and his energy to continue working went down substantially. The family came under excess financial pressure. Children's education is negatively affected. The family became financially indebted. The family was fearful of stigma; therefore, they did not tell others about the disease for long time.

Poverty and Cancer

- Hushed lived in a small village where no good healthcare services were available. In the initial days of pain and suffering, he took pain relief tablets over the counter. It only wasted crucial time of diagnosis. He was diagnosed with cancer at an advanced stage. His treatment started, but the cost was unbearable. He sold inherited land and household items in the process of treatment.
- Later, when his neighbours and community members came to know about his cancer. They came forward and helped him by crowdfunding. He felt bad about losing his physical ability and becoming dependent on others.

fear and family support

- Suman (Name changed), a 26-year-old woman living in Bokaro (a city), Jharkhand, was diagnosed with reproductive cancer. She had been suffering from severe abdominal pain, nausea, and vomiting. Her family took her to the nearest hospital, but the doctors could not determine the disease properly. After continual suffering, she was referred to the RIMS Ranchi for diagnosis. It was eventually diagnosed as reproductive cancer after multiple hospital visits and tests. Suman lived a healthy and satisfied life until she was diagnosed with cancer. She was unable to come to terms with the cancer she was diagnosed with. It took a toll on the poor mental condition of her. She felt completely cut off from the family and community.

fear and family support

- Suman was scared of going through chemotherapy and other treatments and was confused about the future and how to cope with the disease. She felt helpless as she could not do anything to eliminate the disease. Mainly, the fear of death was overwhelming her. The physical pain and fatigue further worsened her mental trauma. She was constantly feeling tired and had no energy to do anything. She also felt isolated as she could not share her pain and suffering with anyone.
- Her parent's struggle to manage treatment expenses was also distressing her. Fortunately, her parents found support from a non-governmental organisation (NGO). The NGO helped them financially and managed counselling sessions for Suman.
- Suman was able to overcome the mental distress and gained confidence to fight the disease.

Stigma and social support

- Nisha (Name changed), a 37-year-old married woman living in Hatia (a rural area), Jharkhand, was diagnosed with colon cancer. She was a mother of seven children. Her husband worked as a watchman. The family belonged to a lower-income stratum.
- She suffered from abdominal pain and sought treatment from local healthcare providers. They could not detect cancer in many consultations. A physician asked to take a cancer test. Then, she was diagnosed with colon cancer.
- Her husband showed confidence that he would do everything to make her free from cancer. Initially, they did not know much about where to go and how to seek treatment. They lost a lot of money and sold assets in private physician treatment, but they got no relief. The family came under huge debt.

Stigma and social support

- Relatives did not pay any attention to the family's distress. Later, her husband isolated her from the rest of the family due to the fear that she could infect others. Her family developed an understanding that this disease is caused due to her past sins. Nisha was told that she had brought shame to the family. As a result, she developed mental health issues like anxiety and insomnia.
- She was brought to Ranchi Cancer Hospital and Research Centre while she went through all the stigma and trauma.

Discussion & Conclusion

Poverty increases with cancer

Illiteracy and lack of awareness make cancer worse

Stigma and apathy in cancer make the problem extremely serious

Discussion & Conclusion

- Cancer and financial distress are connected, and they interplay with stigma and deteriorated mental health. The adverse impact of cancer on poor families in India is phenomenal. It encompasses a lack of resources, fear, and stigma. The lack of public cancer care services also contributes to the problem. In rural and remote areas, patients take a long route to reach cost-effective and quality cancer care facilities.
- Cancer treatments are costly, and all patients cannot afford the necessary care. This can lead to further emotional distress and a more significant burden on the family (Sun et al., 2018; Skelton et al., 2020; de Oliveira et al., 2018; Abdul-Khalek et al., 2020).
- A study found financial difficulties arose due to the cost of treatment and the need to travel to receive it (Jacob et al., 2019). The study argues that patients suffering from cancer stay away from household work due to stigma and fear. This study addressed the NGOs assisting cancer patients in accessing adequate treatment. Similarly, a study argues that the role of philanthropic organisations was providing information about available cancer care facilities, helping them navigate the healthcare system, and sometimes even providing transportation to hospitals and financial aid (Nair et al., 2014).

Discussion & Conclusion

- Cancer patients from Jharkhand frequently face discrimination in receiving health care due to their economic condition. As a result, cancer patients often feel hopeless and isolated, leading to depression and anxiety (Srivastava et al., 2016).
- The challenges posed by poverty are multifaceted and have far-reaching consequences for individuals as well as communities to cope with cancer in this region. Poverty not only limits access to early detection and timely treatment but also exacerbates the physical and emotional burdens faced by cancer patients and their families.
- The role of family and society is very important for a cancer survivor. Psychological support from the family gives hope to life of the cancer patients.
- Healthcare institutions should take counselling services and palliative care.
- In addressing the complex relationship between financial burden and cancer care in Jharkhand, it is the responsibility of both government agencies and non-governmental organisations to implement comprehensive strategies.

References

- Abdul-Khalek, R. A., Guo, P., Sharp, F., Gheorghe, A., Shamieh, O., Kutluk, T., ... & Sullivan, R. (2020). The economic burden of cancer care for Syrian refugees: a population-based modelling study. *The Lancet Oncology*, 21(5), 637-644.
- Alzu'Bi, D., Al-Badarneh, A., & Irshaidat, S. (2022, June). Poverty and Cancer Disproportion in Low-Income Countries. In *2022 13th International Conference on Information and Communication Systems (ICICS)* (pp. 367-372). IEEE.
- Broom, A., & Doron, A. (2012). The rise of cancer in urban India: Cultural understandings, structural inequalities and the emergence of the clinic. *Health*: 16(3), 250-266.
- Callahan, C., & Brintzenhofesoc, K. (2015). Financial quality of life for patients with cancer: An exploratory study. *Journal of Psychosocial Oncology*, 33(4), 377-394.
- Das, S., & Patro, K. C. (2010). Cancer care in the rural areas of India: a firsthand experience of a clinical oncologist and review of literature. *Journal of cancer research and therapeutics*, 6(3), 299-303.
- de Oliveira, C., Weir, S., Rangrej, J., Krahn, M. D., Mittmann, N., Hoch, J. S., ... & Peacock, S. (2018). The economic burden of cancer care in Canada: a population-based cost study. *Canadian Medical Association Open Access Journal*, 6(1), E1-E10.
- Dhillon, P. K., Mathur, P., Nandakumar, A., Fitzmaurice, C., Kumar, G. A., Mehrotra, R., ... & Dandona, L. (2018). The burden of cancers and their variations across the states of India: the Global Burden of Disease Study 1990–2016. *The Lancet Oncology*, 19(10), 1289-1306.
- Ikeda, A., Kawachi, I., Iso, H., Iwasaki, M., Inoue, M., & Tsugane, S. (2013). Social support and cancer incidence and mortality: the JPHC study cohort II. *Cancer Causes & Control*, 24, 847-860.
- Mallath, M. K., Taylor, D. G., Badwe, R. A., Rath, G. K., Shanta, V., Pramesh, C. S., ... & Sullivan, R. (2014). The growing burden of cancer in India: epidemiology and social context. *The Lancet Oncology*, 15(6), e205-e212.
- Nair, M., Parukkutty, K., & Kommadath, S. (2014). Effect of a new social support program by voluntary organization in pediatric oncology department in a developing country. *Pediatric Hematology and Oncology*, 31(3), 212-216.
- Pramesh, C. S., Badwe, R. A., Borthakur, B. B., Chandra, M., Raj, E. H., Kannan, T., ... & Sullivan, R. (2014). Delivery of affordable and equitable cancer care in India. *The Lancet Oncology*, 15(6), e223-e233.
- Rajpal, S., Kumar, A., & Joe, W. (2018). Economic burden of cancer in India: Evidence from cross-sectional nationally representative household survey, 2014. *PLoS one*, 13(2), e0193320.
- Ratcliff, C., Thyle, A., Duomai, S., & Manak, M. (2017). Poverty reduction in India through palliative care: A pilot project. *Indian journal of palliative care*, 23(1), 41.
- Skelton, M., Alameddine, R., Saifi, O., Hammoud, M., Zorkot, M., Daher, M., ... & Mukherji, D. (2020). High-cost cancer treatment across borders in conflict zones: experience of Iraqi patients in Lebanon. *JCO global oncology*, 6, 59-66.
- Srivastava, V., Ansari, M. A., Kumar, A., Shah, A. G., Meena, R. K., Sevach, P., & Singh, O. P. (2016). Study of anxiety and depression among breast cancer patients from North India. *Clinical Psychiatry*, 2(1), 4.
- Sun, L., Legood, R., dos-Santos-Silva, I., Gaiha, S. M., & Sadique, Z. (2018). Global treatment costs of breast cancer by stage: a systematic review. *PLoS one*, 13(11), e0207993.
- Wilkes, G., Freeman, H., & Prout, M. (1994, May). Cancer and poverty: breaking the cycle. In *Seminars in oncology nursing* (Vol. 10, No. 2, pp. 79-88).

Thank you



Imteyaz Ahmad



imteyazahmad777@gmail.com